

Fast Lagrangian Particle Tracking In Unstructured Ocean Model Grids

- Ross Vennell – Cawthron Institute, Nelson, New Zealand



Ocean Dynamics
<https://doi.org/10.1007/s10236-020-01436-7>



Fast lagrangian particle tracking in unstructured ocean model grids

Ross Vennell¹  · Max Scheel¹ · Simon Weppe² · Ben Knight¹ · Malcolm Smeaton¹


Received: 6 May 2020 / Accepted: 23 December 2020
© Springer-Verlag GmbH Germany, part of Springer Nature 2021

¹ Cawthron Institute, Nelson, New Zealand

² MetOcean Solutions, Raglan, New Zealand

Unstructured grids are great!

Achieve resolutions not possible with regular grids




Post model run particle tracking?

- Regular grid – Fast
- Unstructured grids can be very slow!

A day to run 100,000 Particles!


Why speed matters in particle tracking!

Enables tracking millions of particles



- Better statistics:
 - Many release locations and times
 - Heat maps
 - Low probabilities, high significance (e.g. invasive species)
- Many individual behaviours (e.g. parameters drawn from distributions)
- My impatience!

Why is unstructured grid particle tracking so slow?




	SciPy-2D	OT-2D
	90k	90k
Total run time	2190 _{100%}	36 _{100%}
Interpolation total	2179 _{99%}	29 _{79%}


Spatial interpolation velocity, depth, tide, temperature

Three steps:

1. Find triangle containing particle
2. Calculate interpolation weights
3. Apply weights



How much faster than OpenDrift?




200-500 times faster

More importantly... as fast as regular grid!


Finding cell containing particle

- Regular grid - fast by rounding
- Triangular grid was slow, now similar speed!



How?

1) Finding the triangle



The Triangle walk

OpenDrift (Python's SciPy) - poor initial guess

Obvious? - Start at triangle containing particle at last time step


Short Triangle Walk

Average walk is typically 0.7 triangles!

In 3D Schism S-layer model

- *Short Vertical Walk* as well

How? 2) Store and reused interpolation weights



Calculate weights once- use many times!

e.g. time interpolation requires two spatial interpolations at same locations



How? 3) Python Fast?

use Numba: Python loops at C speeds



```
@njit
def add_to(x1,x2,scale, active):
    # x1 += x2*scale for active particles,

    for n in active:
        |   for m in range(x1.shape[1]):
        |       x1[n, m] += x2[n,m]*scale
```



Cook Strait- OpenDrift Comparison





Multi Core performance:



Example: A Biosecurity eDNA inverse problem




B) *Sabella spalanzanii* (combined)




Why speed matters!


Ocean Plastics , public engagement tool



www.oceanplasticsimulator.nz



Statistics on the fly.



Sources of green lipped mussel spat larvae:


120 million particles released over 10 year Schism hindcast,
5 hour run time on 18 cores

➔ Tera Bytes of particle tracks to sort through!

Build heat maps as you go, i.e. don't record
tracks

only mega bytes of output !

OceanTracker



```
params = {  
    'output_folder': 'output',  
    'run_name': 'demo1',  
    'output_file_base': 'demo1_particle',  
    'backtracking': True,  
    'reader': {'field_map': {'water_velocity': ['east_vel', 'north_vel']},  
              'dimension_map': {'space': 'node', 'time': 'time'},  
              'grid_map': {'time': 'time_sec', 'xy': ['east', 'north'], 'triangles': 'tri', 'dry_cells': 'dry_cell_flag'},  
              'time_buffer_size': 200,  
              'file_mask': 'demoHindcast.nc', 'isodate_of_hindcast_time_zero': '2017-01-01'},  
    'particle_release_groups': [{'points': [[1594500, 5482700], [1598000, 5486100]], 'pulse_size': 10,  
                                'release_interval': 3 * 3600},  
                               {'class_name': 'oceantracker.particle_release.polygonRelease.PolygonRelease',  
                                'points': [[1597682.1237, 5489972.7479],  
                                           [1598604.1667, 5490275.5488],  
                                           [1598886.4247, 5489464.0424],  
                                           [1597917.3387, 5489000],  
                                           [1597300, 5489000]]  
                                },  
                               ], 'pulse_size': 10, 'release_interval': 3*3600}],  
    'dispersion': {'A_H': 0.1},  
    'solver': {'n_sub_steps': 6},  
    'user_particle_properties': [{'class_name': 'oceantracker.particle_properties.ageDecay.AgeDecay'},  
                                ]  
}
```

Wrap up.

OceanTracker

- 200-500 times faster than some existing codes
- As fast as regular grid particle tracking
- Enables scaling to millions of particles in 2D and 3D
- Smart- extensible framework for particle tracking
- Inbuilt support for Schism output:
 - works with new LSC vertical grid
 - reads sets of NetCDF files as blocks in read buffer
 - dry cell stranding...



Fast lagrangian particle tracking in unstructured ocean model grids

Ross Vennell¹ · Max Scheel¹ · Simon Weppe² · Ben Knight¹ · Malcolm Smeaton¹

Received: 6 May 2020 / Accepted: 23 December 2020
© Springer-Verlag GmbH Germany, part of Springer Nature 2021

¹ Cawthron Institute, Nelson, New Zealand

² MetOcean Solutions, Raglan, New Zealand